

Quantum Spintronics Design (NV centers in diamond)

Eisuke Abe

RIKEN Center for Quantum Computing

2021.9.8

CMD Spintronics Design Course
(Online)




Diamond envy

III (13)	IV (14)	V (15)
B	C	N



©GIA




$$\rho_N = 1.77 \times 10^{23} \text{ cm}^{-3}$$

Diamond envy

III (13)	IV (14)	V (15)
B	C	N



©GIA




$$\rho_N = 1.77 \times 10^{23} \text{ cm}^{-3}$$

Diamond NV

III (13)	IV (14)	V (15)
B	C	N



©GIA



$$\rho_N = 1.77 \times 10^{23} \text{ cm}^{-3}$$

Diamond NV

Synthetic (CVD) diamond

$2^2 \times 0.5 \text{ mm}^3$, \$700 (E6)


[N] < 5 ppb, [NV] < 0.03 ppb



Not like...




©GIA




$$\rho_N = 1.77 \times 10^{23} \text{ cm}^{-3}$$

Quantum sensing with NV centers

- $B, E, T, S\dots$
- DC & AC modes
- Wide temperature range
- Nondestructive
- High spatial resolution
- Various modalities




Near-surface NV center & NMR



Science 351, 836 (2016)

Scanning probe & condensed matter



Rev. Sci. Instrum. 87, 063703 (2016); Nature 549, 252 (2017)

Outline





- **Basics of NV centers in diamond**
 - Structure
 - Optical & magnetic properties
- **Basics of magnetic resonance**
- **Quantum sensing**
 - AC magnetometry
 - Detection of proton spin ensemble
 - Detection and control of a single proton spin

Outline


- **Basics of NV centers in diamond**
 - Structure
 - Optical & magnetic properties
- **Basics of magnetic resonance**
- **Quantum sensing**
 - AC magnetometry
 - Detection of proton spin ensemble
 - Detection and control of a single proton spin

Crystal & energy level structures

- Negatively-charged (NV^-)
- 4 sp^3 orbitals, 6 e^- (5 from the defect, 1 captured)
- C_{3v} (symmetry axis = quantization axis)





Effective spin-1 system
(e^2 -hole spin-triplet)





Energy levels

C.B. ($E_g = 5.47 \text{ eV} = 227 \text{ nm}$)






PL spectroscopy & imaging

C.B. ($E_g = 5.47 \text{ eV} = 227 \text{ nm}$)




PL spectroscopy & imaging

C.B. ($E_g = 5.47 \text{ eV} = 227 \text{ nm}$)




Photon statistics

C.B. ($E_g = 5.47 \text{ eV} = 227 \text{ nm}$)




One photon at a time




V.B.

Time-resolved fluorescence

C.B. ($E_g = 5.47 \text{ eV} = 227 \text{ nm}$)




The **non-radiative & spin-selective** channel provides a means to **read out & initialize** the NV spin



Optically detected magnetic resonance


C.B. ($E_g = 5.47 \text{ eV} = 227 \text{ nm}$)




$$H = DS_z^2 + \gamma_e B_0 S_z$$

$D = 2.87 \text{ GHz}, \gamma_e = 28 \text{ MHz/mT}$


$B_0 = 4.7 \text{ mT}, f_{\pm} = 2.87 \pm 0.132 \text{ GHz}$





Experimental setup



Experimental setup



Experimental setup



AIP Adv. **10**, 025206 (2020)
Misonou *et al.*

Outline

- **Basics of NV centers in diamond**
 - Structure
 - Optical & magnetic properties
- **Basics of magnetic resonance**
- **Quantum sensing**
 - AC magnetometry
 - Detection of proton spin ensemble
 - Detection and control of a single proton spin

Magnetic resonance

Torque equation (Larmor precession)


$$\frac{d\mu}{dt} = \mu \times \gamma B_0$$

↑
↑
Gyromagnetic ratio

Magnetic moment: $\mu = \gamma J$




(from Wikipedia)




Frame rotating at angular velocity Ω :
Rotate slower...why?

Joseph Larmor
(1857–1942)

Magnetic resonance



AC field rotating in the xy plane at Ω




Frame rotating at angular velocity Ω :

Rotate slower...why?


DC field along the z direction becomes weaker

Magnetic resonance

Frame rotating at $\Omega = \gamma B_0$



Rest (non-resonant) frame




- Rotations about the $\pm \hat{x}$, $\pm \hat{y}$ axes are realized by adjusting the microwave phases
- Rotation about the \hat{z} axis is superposed when observed from the rest (non-resonant) frame




Qubit & Bloch sphere

Qubit, spin-½ (NV is spin-1!)


$$\begin{cases} |"0"\rangle \equiv |m_s = 0\rangle \\ |"1"\rangle \equiv |m_s = -1\rangle \end{cases}$$



Rabi oscillation



Quantum coherence



$$|0\rangle \equiv |m_s = 0\rangle$$


$$|\Psi\rangle = \alpha|0\rangle + \beta|1\rangle$$

$$|1\rangle \equiv |m_s = -1\rangle$$


T_2 : measure of how long a superposition state is preserved

Measurement of T_2

Time domain → Decay time constant T_2




Frequency domain → Lorentz width




Measurement of T_2

Time domain → Decay time constant T_2





Frequency domain → Lorentz width




Inhomogeneous line broadening


Seen to precess at $\delta\nu$ in the frame rotating at ν_e



Spin echo




Modulation function




Phase accumulation

by DC field


(angle from the y axis)




$\pi/2$




$t = 0$




$t = \tau^{(-)}$




$t = \tau^{(+)}$



$t = 2\tau$



Coherence time



Stretched exponential decay

$$\exp \left[- \left(\frac{2\tau}{T_2} \right)^p \right]$$



Near-surface NV center

- N^+ implantation into ^{12}C ($I = 0$) layer
- $d_{\text{NV}} = 6.26 \text{ nm}$
- $B_0 = 23.5 \text{ mT}$

Outline

- **Basics of NV centers in diamond**
 - Structure
 - Optical & magnetic properties
- **Basics of magnetic resonance**
- **Quantum sensing**
 - AC magnetometry
 - Detection of proton spin ensemble
 - Detection and control of a single proton spin

Nuclear spin sensing




Nuclear spins **precess** at $f_{\text{ac}} = \text{a few kHz--MHz}$ under B_0

→ Weak AC magnetic field on the NV spin


→ Detect using quantum coherence

AC magnetometry

CP ($N = 4$)




Modulation function




AC field at $f_{ac} = 1/2\tau$

Sensor phase buildup = loss of coherence




AC magnetometry

CP ($N = 4$)




Modulation function




- φ depends on the **initial phase α of the AC field** ($\varphi \propto \cos \alpha$)
- Average over **random α**


Nuclear spin sensing



- $T_2 = 6.2 \mu\text{s} @ B_0 = 23.5 \text{ mT}$
- $N = 64$ (XY16)
- $2\tau = 2 \times 32 \mu\text{s}/64 = 1 \mu\text{s}$
 $\rightarrow \gamma_H B_0 = (42.577 \text{ kHz/mT}) \times B_0 = 1.00 \text{ MHz}$




Nuclear spin sensing



$$C(\tau) = f(B_{\text{rms}})$$

$$B_{\text{rms}} = \frac{\mu_0}{4\pi} h \gamma_H \sqrt{\frac{5\pi\rho}{96d_{\text{NV}}^3}}$$


Phys. Rev. B **93**, 045425 (2016)




- Proton density $\rho = 6 \times 10^{28} \text{ m}^{-3}$ (known)
- $d_{\text{NV}} = 6.26 \text{ nm}$
- $B_{\text{rms}} \approx 560 \text{ nT}$
- Detection volume $(d_{\text{NV}})^3 \approx 0.25 \text{ zL}$ (zepto = 10^{-21})
- # of proton $\rho(d_{\text{NV}})^3 \approx 1500$
- Thermal pol. (10^{-7}) vs. statistical pol. $(1500)^{0.5} \approx 39$

Toward single-molecular imaging


- **Strategy**
 - Detect **individual nuclear spins** in a single molecule
 - Determine their **nuclear species (& chemical shifts) and positions**
- **Practical issues**
 - Preparation of high-quality near-surface NV centers
 - Accurate positioning of single molecules/proteins near the sensor



Detection of single protons in CVD diamond
(as a testbed)




Nuclear spin sensing



- Single NV in a N-doped CVD film ($[{}^{12}\text{C}] = 99.999\%$)
- $N = 64$ (XY16)
- $f_H = \gamma_H B_0 = 42.577 \text{ kHz/mT} \times 28.7 \text{ mT} = 1.2239 \text{ MHz}$

Correlation spectroscopy



The transition probability for random phases (α)

→ Sweep t_{corr}

$$p(t_1) \approx \frac{1}{2} \left\{ 1 - \frac{1}{2} \left(\frac{\gamma B_{\text{ac}} t_s}{\pi} \right)^2 \cos(2\pi f_{\text{ac}} t_{\text{corr}}) \right\}$$



Nature Commun. **4**, 1651 (2013) Laraoui *et al.*

Phys. Rev. Appl. **4**, 024004 (2015) Kong *et al.*

Nature Commun. **6**, 8527 (2015) Staudacher *et al.*


Phys. Rev. Lett. **116**, 197601 (2016) Boss *et al.*

Correlation spectroscopy of a nucleus



- $f_0 = 1.2234 \text{ MHz}$
- $f_1 = 1.2046 \text{ MHz}$


Correlation spectroscopy of a nucleus



Hamiltonian of NV-¹H coupled system

$$H = f_H I_z + |m_s = -1\rangle \langle -1| (A_{\parallel} I_z + A_{\perp} I_x)$$


→ No hyperfine field when $|m_s = 0\rangle$



- $f_0 = 1.2234$ MHz = f_H ($m_s = 0$)
- $f_1 = 1.2046$ MHz = $f_H + A'_{\parallel}$ ($m_s = -1$)

$A'_{\parallel} = -18.8$ kHz
 $(f_0 + f_1)/2 = 1.2140$ MHz → dip


Coherent control of a nuclear spin



Hamiltonian of NV-¹H coupled system


$$H = f_H I_z + |m_s = -1\rangle \langle -1| (A_{\parallel} I_z + A_{\perp} I_x)$$

→ The single ¹H *n*-spin rotates about the A_{\perp} axis





- $N \rightarrow 656$ ($\tau = 412.5$ ns, fixed)
- $f_{\text{osc}} = 7.414$ kHz = $A'_{\perp}/2$


Conditional rotation of a nuclear spin




Evolution of n -spin vector




Start from $|m_s = 0\rangle$



Start from $|m_s = -1\rangle$




Coherent control of a nuclear spin



Transition probability of the NV spin

$$P_{0,X} = 1 - \frac{1}{2} \underbrace{(1 - \mathbf{n}_0 \cdot \mathbf{n}_{-1})}_{-1} \sin^2 \frac{N\phi_{cp}}{2}$$

Phys. Rev. Lett. **109**, 137602 (2012) Taminiau *et al.*




- $N \rightarrow 656$ ($\tau = 412.5$ ns, fixed)
- $f_{osc} = 7.414$ kHz = $A'_{\perp}/2$

$P_X < 0.5 \rightarrow$ Single proton!

Appl. Phys. Lett. **117**, 114002 (2020) Sasaki *et al.*

Determination of hyperfine constants



Magnetic dipole interaction

$$A_{\parallel} = h\gamma_e\gamma_H \frac{3 \cos^2 \theta - 1}{r^3}$$

$$A_{\perp} = h\gamma_e\gamma_H \frac{3 \cos \theta \sin \theta}{r^3}$$

$$\begin{aligned} A_{\parallel} &= -19.0 \text{ kHz} \\ A_{\perp} &= 22.9 \text{ kHz} \end{aligned}$$




$$\begin{aligned} r &= 1.44 \text{ nm} \\ \theta &= 72.3^\circ \end{aligned}$$

The position of the nucleus can be determined
→ Molecular structure analysis

(Azimuthal angle ϕ) Phys. Rev. B **98**, 121405 (2018) Sasaki *et al.*

Appl. Phys. Lett. **117**, 114002 (2020) Sasaki *et al.*

Magnetic field dependence




($\gamma_H/\gamma_C = 3.97 \rightarrow$ Spurious harmonics?) Phys. Rev. X 5, 021009 (2015) Loretz *et al.*


Appl. Phys. Lett. 117, 114002 (2020) Sasaki *et al.*

Toward single-molecular imaging

- **Spatial resolution**
 - Accurate measurement of $e-n$ int. const's (A_{\parallel}, A_{\perp}) $\approx (r, \theta)$
 - Information on the azimuthal angle ϕ
- **Spectral resolution**
 - Nuclear species
 - J -couplings & chemical shifts (ppm!)
 - Limited by sensor/memory lifetimes ($T_{2e/n}, T_{1e/n}$)




AC magnetometry revisited




- φ depends on the **initial phase α of the AC field** ($\varphi \propto \cos \alpha$)
- Average over **random α**

Ultrahigh resolution sensing






- φ depends on the **initial phase α of the AC field** ($\varphi \propto \cos \alpha$)
- Average over **random α**
- **If the data acq. is periodic**, adjacent α 's are related by $\alpha_{k+1} = 2\pi f_{\text{ac}} t_L + \alpha_k$

Ultrahigh resolution sensing




Ultrahigh resolution sensing

$B_{\text{ac}} = 96.5 \text{ nT}$ & $f_{\text{ac}} = 2.001 \text{ MHz}$ applied from a coil, detected by a single NV center




- $T = 3600 \text{ sec}$
- $\text{FWHM} = 0.304 \text{ mHz}$

Free induction decay



Nuclear Rabi oscillation



- $T_{\text{rf}, \pi/2} = 4.115 \mu\text{s}$
- [PulsePol]– $T_{\text{rf}, \pi/2}$ –[X/2–(XY16)–Y/2–L_{RO}]⁵⁰
- $f_{\text{sample}} = 1/t_L = 84.46 \text{ kHz}$
- $f_p = (f_0 + f_1)(t_s/t_L)/2 + f_0(t_s - t_L)/t_L = 1.2182 \text{ MHz}$
→ Split (analogous to chemical shifts)

Summary

- **Tools for single-molecule imaging/structural analysis are being developed**
 - Ultrahigh resolution sensing^[1,2,3], resolving chemical shifts^[3,4] & suppression of back action from n -spins^[5,6]
 - Determination of the position of individual n -spins^[7,8,9]
 - Detection and control of single proton spins^[10,11]

- [1] Science **356**, 832 (2017) Schmitt *et al.* (Ulm)
- [2] Science **356**, 837 (2017) Boss *et al.* (ETH)
- [3] Nature **555**, 351 (2018) Glenn *et al.* (Harvard)
- [4] Science **357**, 67 (2017) Aslam *et al.* (Stuttgart)
- [5] Nature Commun. **10**, 594 (2019) Pfender *et al.* (Stuttgart)
- [6] Nature **571**, 230 (2019) Cujia *et al.* (ETH)
- [7] Phys. Rev. B **98**, 121405 (2018) Sasaki *et al.* (Keio)
- [8] Phys. Rev. Lett. **121**, 170801 (2018) Zopes *et al.* (ETH)
- [9] Nature **576**, 411 (2019) Abobeih *et al.* (Delft)
- [10] Phys. Rev. Lett. **113**, 197601 (2014) Sushkov *et al.* (Harvard)
- [11] Appl. Phys. Lett. **117**, 114002 (2020) Sasaki *et al.* (Keio)